

Calculation of thermal conductivity coefficients for magnetized neutron star

M. V. Glushikhina* and G. S. Bisnovatyi-Kogan**

Space Research Institute of Russian Academy of Sciences, Moscow, Russia

*E-mail: m.glushikhina@iki.rssi.ru

**E-mail: gkogan@iki.rssi.ru

Abstract

The coefficients that determine the electron heat transfer and diffusion in the crust of neutron stars are calculated on the basis of a solution of the Boltzmann equation with allowance for degeneracy.

1 Introduction

Thermal conductivity in the envelopes of neutron stars plays crucial role in many aspects of evolution of these stars. Thermal conductivity is the basic quantity needed for calculating the relationship between the internal temperature of a neutron star and its effective surface temperature, this relationship affects thermal evolution of the neutron star and its radiation spectra. We should know the transport properties of dense matter where electrons are strongly degenerate and form nearly ideal Fermi-gas. Under such conditions, electrons are usually most important heat carriers. The present work is devoted to calculation of electron thermal conductivity coefficients under the given conditions on the basis of the solution of Boltzmann equation. The magnetic field limits the motion of electrons in the direction perpendicular to the field lines and, since they are the main carriers of the heat transport, the thermal conductivity in this direction is highly suppressed, while remaining unaffected along the field lines. In the present work we obtain a new approximation for coefficients of electron heat conductivity along and across magnetic field lines.

2 Boltzmann equations and transfer equations

To calculate the transport coefficients we use a Boltzmann equation with allowance for degeneracy and only two-body collisions. We consider an electron gas in the crystal lattice of heavy nuclei and we take into account the interaction of the electrons with the nondegenerate nuclei and with one another. The nuclear component of the matter in the crust is evidently in the crystal state and therefore the isotropic part of the distribution function f_{N0} may differ from the Maxwellian distribution. If the mass of the m_N nucleus is much greater, than the electron mass m_e , the to the terms $\sim m_e/m_N$ the details of the distribution function f_{N0} is unimportant, and the calculations can be made for arbitrary f_{N0} . The transfer equations for the electron concentration, total momentum and energy in the two-component mixture of electrons and nuclei can be obtained in the usual manner [1] from the Boltzmann equation for nonrelativistic electrons. [1–4]

$$\frac{dn_e}{dt} + n_e \frac{\partial c_{0i}}{\partial r_i} + \frac{\partial}{\partial r_i} (n_e \langle v_i \rangle) = 0 \quad (1)$$

$$\rho_e \frac{dc_{0i}}{dt} = \frac{1}{c} \epsilon_{ikl} j_k B_l - \frac{\partial \Pi_{ik}}{\partial r_i} + \varepsilon (E_i + \frac{1}{c} \epsilon_{ikl} c_{0k} B_l) \quad (2)$$

$$\frac{3}{2} k n_e \frac{dT}{dt} - \frac{3}{2} k T \frac{\partial}{\partial r_i} (n_e \langle v_i \rangle) + \frac{\partial q_{ei}}{\partial r_i} + \Pi_{ik} \frac{\partial c_{0i}}{\partial r_i} = j_i (E_i + \frac{1}{c} \epsilon_{ikl} c_{0k} B_l) - \rho_e \langle v_i \rangle \frac{dc_{0i}}{dt} \quad (3)$$

Where:

$$\begin{aligned} p_e &= \frac{1}{3} n_e m_e \langle v^2 \rangle & \Pi_{ik} &= n m \langle v_i v_k \rangle \\ q_i &= \frac{1}{2} n m \langle v^2 v_i \rangle & \langle v_i \rangle &= \frac{1}{n} \int f v_i d c_i \end{aligned} \quad (4)$$

c_{0i} is mass-average velocity, v_i is the thermal velocity of the electrons.

3 Derivation of general expressions for the transport coefficients

The Boltzmann equation can be solved by the Chapmen-Enskog method of successive approximation [1]. The zeroth approximation to the electron distribution function is found by equating to zero the collision integral:

$$f_0 = [1 + \exp \frac{m_e v^2 - 2\mu}{2kT}]^{-1} \quad n_e = D \int f_0 dv_i \quad (5)$$

Here, μ is chemical potential of electrons, k is Boltzmann's constant, T is the temperature, and $D = 2m^3/(2\pi\hbar)^3$.

The nuclear distribution function in the zeroth approximation f_{N0} is assumed to be isotropic with respect to the velocities and to depend on the local thermodynamic parameters; otherwise it can be arbitrary with the normalization:

$$n_N = \int f_{N0} dc_{Ni}$$

Using (5) in (1)-(4), we obtain the zeroth approximation for the transfer equations. In this approximation $\langle v_i \rangle = 0$, $q_i = 0$, $\Pi_{ik} = (\Pi_e + \Pi_N)\delta_{ik}$

$$n_e = 2\left(\frac{kTm_e}{2\pi^2\hbar^2}\right)^{3/2} G_{3/2}(x_0) \quad P_e = 2kT\left(\frac{kTm_e}{2\pi^2\hbar^2}\right)^{3/2} G_{5/2}(x_0) \quad (6)$$

$$G_n(x_0) = \frac{1}{\Gamma(n)} \int \frac{x^{n-1} dx}{1 + \exp(x - x_0)} \quad x_0 = \frac{\mu}{kT}$$

In what follows, instead of $G_n(x_0)$ we will write G_n cause the argument is the same. In the first approximation, we seek the function f in the form:

$$f = f_0(1 + \chi(1 - f_0)) \quad (7)$$

We take deviation of the nuclear distribution function from zeroth approximation in the form:

$$f_N = f_{N0}(1 + \chi_N) \quad (8)$$

χ is linear and admits representation of the solution in the form:

$$\chi = A_i \frac{\partial \ln T}{\partial r_i} - n_e D_i d_i \frac{G_{5/2}}{G_{3/2}} \quad \chi_N = -A_{Ni} \frac{\partial \ln T}{\partial r_i} - n_e D_{Ni} d_i \frac{G_{5/2}}{G_{3/2}} \quad (9)$$

The functions A_i, A_{Ni} and D_i, D_{Ni} determine diffusion and heat transfer. Substituting (9) in the equation for χ we can obtain equations for A_i, A_{Ni}, D_i, D_{Ni} [6]

$$\begin{cases} f_0(1 - f_0)\left(\frac{m_e v^2}{2kT} - \frac{5G_{5/2}}{2G_{3/2}}\right)v_i = I_{ee}(A_i) + I_{eN}(A_i) \\ \frac{1}{n_e} f_0(1 - f_0)v_i = I_{ee}(D_i) + I_{eN}(D_i) \end{cases} \quad (10)$$

We seek solution of (10) in the form of an expansion in polynomials Q_n that are orthogonal with weight $f_0(1 - f_0)x^{3/2}$. Where Q_n - are analogous to Sonine polynomials [5].

$$\begin{aligned} Q_0(x) &= 1 & x &= u^2 = \frac{m_e}{2kT} v_i^2 \\ Q_1(x) &= \frac{5G_{5/2}}{2G_{3/2}} - x \end{aligned} \quad (11)$$

We seek A_i and D_i in the form:

$$A_i = (a_0 Q_0 + a_1 Q_1)v_i \quad D_i = (d_0 Q_0 + d_1 Q_1)v_i \quad (12)$$

Multiplying (10) by $DQ_0(x)u_i$ and $DQ_1(x)u_i$ and integrating with respect to dc_i , we obtain a system of equation for heat conductivity coefficients:

$$\begin{cases} 0 = 3i \frac{eBn_e}{m_e c} a_0 + a_0(a_{00} + b_{00}) + a_1(a_{01} + b_{01}) \\ -\frac{15}{4} n_e \left(\frac{7G_{7/2}}{2G_{5/2}} - \frac{5G_{5/2}^2}{2G_{3/2}^2} \right) = i \frac{15eBn_e}{4m_e c} a_1 + a_0(a_{10} + b_{10}) + a_1(a_{11} + b_{11}) \end{cases} \quad (13)$$

Here a_{jk} b_{jk} are matrix elements for collision integrals.

4 Finding of matrix elements

$$a_{jk} = D^2 \int f_0 f_{01} (1 - f'_0) (1 - f'_{01}) Q_j(u^2) u_i [Q_k(u^2) + Q_k(u_1^2) u_{1i} - Q_k(u'^2) u'_i - Q_k(u_1'^2 u_{1i}')] g_{ee} W_{ee}(\theta, g_{ee}) d\Omega dc_{1i} dc_i \quad (14)$$

$$b_{0j} = D \int f_0 f_{N0} (1 - f'_0) Q_j(u^2) u_i [u_i - u'_i - \frac{n_e}{n_N} (\frac{m_e}{m_N})^{1/2} (u_{Ni} - u'_{Ni})] g_{eN} W_{eN}(\theta, g_{eN}) d\Omega dc_{Ni} dc_i \quad (15)$$

$$b_{jk} = D \int f_0 f_{N0} (1 - f'_0) Q_j(u^2) u_i [Q_k(u^2) u_i - Q_k(u'^2) u'_i] g_{eN} W_{eN}(\theta, g_{eN}) d\Omega dc_{Ni} dc_i, \quad k \geq 1 \quad (16)$$

$W(\theta, g) d\Omega$ is the effective differential cross section for scattering of particles with relative velocity g that is deflected through angle θ and after the collision lies in the solid angle $d\Omega$.

5 Tensor of heat conductivity

We can find a_0 and a_1 by solving (13) and using following expressions:

$$\begin{aligned} a_0 &= a_0^1 + iBb_0^1 & a_1 &= a_1^1 + iBb_1^1 \\ c_0^1 &= (a_0^1)_{B=0} - a_0^1 & c_1^1 &= (a_1^1)_{B=0} - a_1^1 \end{aligned} \quad (17)$$

we can find coefficients of heat conductivity tensor [3, 4, 8].

$$\lambda_{ik} = \frac{5}{2} \frac{k^2 T n}{m} ((a_0^1 - a_1^1) \delta_{ik} - \epsilon_{ikn} B_n (b_0^1 - b_1^1) + B_i B_k (c_0^1 - c_1^1))$$

In previous works [7, 9] the following approximation was used:

$$\frac{\lambda_{\perp}}{\lambda_{\parallel}} = \frac{1}{1 + (\omega\tau)^2} \quad (18)$$

Considering two cases of strong and weak degeneracy, we obtain the following relations: for weak degeneracy (nondegenerate):

$$\frac{\lambda_{\perp}}{\lambda_{\parallel}} = \frac{1}{1 + \frac{17(\omega\tau)^2 + 2.76(\omega\tau)^4}{2.58 + 0.39(\omega\tau)^2}} \quad (19)$$

for strong degeneracy:

$$\frac{\lambda_{\perp}}{\lambda_{\parallel}} = \frac{1}{1 + \frac{0.31(\omega\tau)^2 + 0.55(\omega\tau)^4}{0.14 + 0.6(\omega\tau)^2}} \quad (20)$$

here in weak degeneracy case $\tau = \frac{3\sqrt{m_e}(kT)^{3/2}}{4\sqrt{2\pi}Z^2e^4n_N\Lambda}$ is relaxation time between electron-nucleus collisions and in

strong degeneracy case $\tau = \frac{3h^3n_e}{32\pi^2Z^2e^4n_Nm_e\Lambda}$ and $\omega = \frac{eB}{m_e c}$ is electron cyclotron frequency. $\lambda_{\parallel} = \frac{k^2 n_e^2 T h^3}{32m^2 Z^2 e^4 n_N \Lambda}$ in the strong degeneracy case and $\lambda_{\parallel} = 4.4k \frac{n_e}{n_N \sqrt{2\pi}} (\frac{kT}{m})^{1/2} \frac{(kT)^2}{Z^2 e^4 \Lambda}$ in the weak degeneracy case (nondegenerate).

In our work we obtain the more exact relation for the heat conductivity coefficients along and across magnetic field lines by solving Boltzmann kinetic equation.

6 Acknowledgements

The work of GSBK and MVG was partially supported by the Russian Foundation for Basic Research grant 11-02-00602 and the Russian Federation President Grant for Support of Leading Scientific Schools NSh-5440.2012.2.

The work of MVG was also partially supported by Russian Federation President Grant for Support of Young Scientists MK-2918.2013.2.

References

- [1] S. Chapmen and T.G. Cowling, Mathematical Theory of Nonuniform Gases, Cambrige(1952)[Russian translation published by IL, Moscow, 1958].
- [2] W. Marshall, The kinetic theory of an ionized gas. p. 3 At. Energy Res Estable 1960, N T/R, 2419.
- [3] G.S. Bisnovaty-Kogan, Transport properties of partially ionized two-temperature plasma and isotropic corrections to maxvillian distribution function (in Russian), Diploma thesis, Moscow Institute of Physics and Technology, 1964.
- [4] G.S. Bisnovaty-Kogan, Heat transfer and diffusion of partially ionized two-temperature plasma, Journal of Applied Mechanics and Technical Physics, No. 3, p. 43, 1964.
- [5] G.S. Bisnovaty-Kogan, M.M. Romanova, Neutron diffusion and heat transfer in the crusts of neutron stars, JETP **56 (2)**, 1982.
- [6] L. Landau, E. Lifshitz, Statisticheskaya fizika, Fizmatlit, Moscow (2002).
- [7] E. Flowers,N. Itoh, Transport properties of dense matter, Ap. J. **206**: 218-242, 1976.
- [8] L.E. Kalikhman, Elements of magnetohydrodynamics, Atomizdat, Moscow (1964).
- [9] D.G. Yakovlev, V.A. Urpin, On thermal and electrical conductivities in neutron stars and white dwarfs, Astron. J. **57**: 526-536, 1980.